

FINAL REPORT  
 "<sup>22</sup>Na Decay Gamma Rays From Classical Novae"  
 NASA NAG5-1565  
 5-15-91 to 5-14-93

James W. Truran  
 Department of Astronomy and Astrophysics  
 University of Chicago  
 Chicago, IL

FINAL  
 IW-93-CR  
 37 CIT.  
 33549  
 P-5

SUMMARY OF RESEARCH ACTIVITIES  
 SUPPORTED IN FULL OR IN PART BY  
 NASA GRANT NAG 5-1565  
 AT THE UNIVERSITY OF ILLINOIS

NASA Grant NAG 5-1565, at the University of Illinois, has provided support for a program of theoretical research in nuclear astrophysics and related areas, focusing upon the possibility of detection of gamma rays from nearby novae. Particular attention has been given to the evaluation of the theoretical expectations for gamma ray emission from four possible sources: (1) the positron decays of the unstable CNO and fluorine isotopes that are transported to the surface regions of the envelope in the earliest stages of the outbursts; (2) <sup>7</sup>Be decay gamma rays; (3) <sup>22</sup>Na decay gamma rays released in the later stages of the outbursts; and (4) <sup>26</sup>Al decay gamma rays from novae and their possible contributions to Galactic emission. The critical questions of (i) the frequency of occurrence of ONeMg-enriched novae; (ii) the expected Galactic distribution of the novae that produce <sup>26</sup>Al; and (iii) the nature of the observed soft X-ray emission from classical novae, have also been addressed. Considerable progress in research has been achieved on many of these fronts. Researchers who have received some level of support from this grant include the Principal Investigator, James W. Truran, Sumner Starrfield, Ami Glasner, Anurag Shankar, and John Hayes. Brief summaries of the results of several research projects are presented below.

Powering the Light Curves of Fast Classical Novae

Observations of fast classical novae indicate the existence of a large and serious disagreement with theory, regarding the maximum luminosities achieved by these systems. Fast novae in our Galaxy, and those in M-31 and the LMC (for which distances are particularly well determined), reach absolute visual magnitudes as bright as -8.5 and can remain brighter than -7 for several days. This behavior has not been successfully reproduced by any self-consistent, time-dependent numerical simulation. Hayes and Truran (1994) have demonstrated that existing numerical models are energetically capable of powering such observed emission for the requisite timescale, and argue that the discrepancy arises due to the fact that the physics describing the exchange of energy and momentum between the material and radiation has been oversimplified in the hydrodynamical studies published to date. Efforts are currently underway to remedy this situation, by substantial modification of the numerical procedures utilized in our hydrodynamic program.

(NASA-CR-197454) Na-22 DECAY GAMMA  
 RAYS FROM CLASSICAL NOVAE Final  
 Report, 15 May 1991 - 14 May 1993  
 (Chicago Univ.) 5 p

N95-16383

Unclass

## Abundances in Nova Ejecta

Livio and Truran (1994) have re-examined the question of the frequency of occurrence of oxygen-neon-magnesium (ONeMg) degenerate dwarfs in classical nova systems, in light of recent observations which have been interpreted as suggesting that “neon novae” can be associated with relatively low mass white dwarfs. Determinations of heavy element concentrations in nova ejecta were reviewed, and possible interpretations of their origin were examined. We concluded that, of the eighteen classical novae for which detailed abundance analyses are available, only two (or possibly three) seem unambiguously to demand the presence of an underlying ONeMg white dwarf: V693 CrA 1981, V1370 Aql 1982, and possibly QU Vul 1984. Three other novae which exhibit significant neon enrichments, relative to their total heavy element concentrations, are RR Pic 1925, V977 Sco 1989, and LMC 1990 #1. This result is entirely consistent with present frequency estimates, and our interpretation of the lower levels of enrichment in other systems explains, in a natural way, the existence of relatively low mass white dwarfs in some of the “neon” nova systems.

## The Nature of the Soft X-Ray Emission from GQ Muscae

Ögelman *et al.* (1993) have recently reported the detection of Nova GQ Muscae 1983 at soft x-ray wavelengths, nine years after outburst, with the ROSAT satellite. The observed spectrum is very soft, and is generally consistent with black body emission from a  $\sim M_{\odot}$  white dwarf, burning at a near Eddington luminosity and an effective temperature  $\sim 3.4 \times 10^5$  K. Such a soft x-ray signature is entirely consistent with expectations for the long term evolution of classical novae in outburst (MacDonald, Fujimoto, and Truran 1985), as it can arise as a natural consequence of the hardening of the radiation from novae, as the photospheric radius decreases during an extended phase of shell hydrogen burning at approximately constant bolometric luminosity. Truran, Glasner, and Hayes (1994), in a paper soon to be submitted for publication: (1) review and discuss the observations of Nova GQ Muscae, the prototype event of this nature, (2) present recent calculations which allow us better to understand the timescales for the onset and duration of the observed phase of soft x-ray emission, and (3) address the obvious question as to why it is that GQ Muscae is one of only a few of the recent classical novae to have been found to exhibit this behavior.

## Nucleosynthesis Associated with ONeMg Enriched Novae

Detailed calculations have been performed of the nucleosynthesis that accompanies the occurrence of thermonuclear outbursts on ONeMg white dwarfs (Politano *et al.* 1995). The trends in the production of the gamma ray sources  $^{23}\text{Na}$  and  $^{26}\text{Al}$ , as a function of the mass of the white dwarf have also been identified. Careful attention is being given to the interpretation of the abundance observations of novae, in lieu of the fact that there appears to be an inconsistency: the large masses of the ejected shells that have been inferred from observations of several recent such “neon novae” are inconsistent with the large masses expected from theory for the ONeMg white dwarfs on which such outbursts are expected to occur. Revised rates for a number of important thermonuclear reactions for  $^{23}\text{Na}$  and  $^{26}\text{Al}$  production have been included, which allow more realistic predictions of the emerging abundances.

PUBLICATIONS  
SUPPORTED IN FULL OR IN PART BY  
NASA GRANTS NAG 5-1565 AND NAG 5-2081  
"Gamma Rays from Classical Novae"  
AT THE UNIVERSITY OF ILLINOIS 1991-1994

1. Starrfield, S., Truran, J.W., Sparks, W.M., and Krautter, J., "Observations and Predictions of EUV Emission from Classical Novae," in Extreme Ultraviolet Astronomy, R. Malina and S. Bowyer (Eds.), Pergamon Press, New York, pp. 168-176, 1991.
2. Starrfield, S., Sparks, W.M., Truran, J.W., and Shaviv, G., "Neon Novae, Recurrent Novae, and Type I Supernovae," in Supernovae, S.E. Woosley (Ed.), Springer-Verlag, New York, pp. 602-605, 1991.
3. Saizar, P., Starrfield, S., Ferland, G.J., Wagner, R.M., Truran, J.W., Kenyon, S.J., Sparks, W.M., Williams, R.E., and Stryker, L.L., "PW Vul: A Nova With Nearly Solar Abundances," Astrophysical Journal, Vol. 367, pp. 310-320, 1991.
4. Shankar, A., Livio, M., and Truran, J.W., "The Common Envelope Phase in Classical Novae: One Dimensional Models," Astrophysical Journal, Vol. 374, pp. 623-630, 1991.
5. Starrfield, S., Politano, M., Truran, J.W., Sparks, W.M., and Weiss, A., "Hydrodynamic Studies of Accretion onto ONeMg White Dwarfs Using a Large Nuclear Reaction Network," Bulletin of the American Astronomical Society, Vol. 23, p. 967, 1991.
6. Politano, M., Starrfield, S., Sparks, W.M., and Truran, J.W., "Hydrodynamic Studies of Accretion onto Massive White Dwarfs: ONeMg-Enriched Nova Outbursts," Bulletin of the American Astronomical Society, Vol. 23, p. 1324, 1991.
7. Brown, L.E., Bazan, G., Clayton, D.D., and Truran, J.W., "Hot Bottom Burning in AGB Stellar Envelopes," Bulletin of the American Astronomical Society, Vol. 23, p. 1400, 1991.
8. Saizar, P., Ferland, G.J., Wagner, R.M., Starrfield, S., Stryker, L., Sparks, W.M., Williams, R.E., Truran, J.W., and Kenyon, S.J., "Multiwavelength Studies of Classical Nova Shells," Bulletin of the American Astronomical Society, Vol. 23, p. 1463, 1991.
9. Truran, J.W., Starrfield, S., and Sparks, W.M., "X-Ray and  $\gamma$ -Ray Emission from Classical Nova Outbursts," in The Compton Observatory Science Workshop, C.R. Shrader, N. Gehrels, and B. Dennis (Eds.), NASA Conference Publications 3137, pp. 369-376, 1992.
10. Starrfield, S., Politano, M., Truran, J.W., and Sparks, W.M., "A Burst from a Thermonuclear Runaway on an ONeMg White Dwarf," in The Compton Observatory Science Workshop, C.R. Shrader, N. Gehrels, and B. Dennis (Eds.), NASA Conference Publications 3137, pp. 337-386, 1992.
11. Livio, M. and Truran, J.W., "Type I Supernovae and Accretion Induced Collapses from Cataclysmic Variables," Astrophysical Journal, Vol. 389, pp. 695-703, 1992.
12. Livio, M., Mostichiadis, A., Ögelman, H., and Truran, J.W., "On the Role of Radioactive Decays in Powering x-Rays from Novae," Astrophysical Journal, Vol. 394, pp. 217-220, 1992.
13. Saizar, P., Starrfield, S., Ferland, G.J., Wagner, R.M., Truran, J.W., Kenyon, S.J., Sparks, W.M., Williams, R.E., and Stryker, L.L., "A Multiwavelength Study of Nova QU Vul 1984," Astrophysical Journal, Vol. 398, pp. 651-664, 1992.

14. Starrfield, S., Shore, S.N., Sparks, W.M., Sonneborn, G., Truran, J.W., and Politano, M., "The  $\gamma$ -Ray Flux from Nova Herculis 1991," Astrophysical Journal Letters, Vol. 391, pp. L71-L74, 1992.
15. Starrfield, S., Truran, J.W., Politano, M., Sparks, W.M., Nofar, I., and Shaviv, G., " $^{22}\text{Na}$  and  $^{26}\text{Al}$  Production in Nova Outbursts," Physics Reports, Vol. 227, pp. 223-234, 1993.
16. Gehrz, R., Truran, J.W., and Williams, R.E., "Novae: Contributions to the Interstellar Medium," in Protostars and Planets III, Levy, E.H. and Lunine, J.I. (Eds.), University of Arizona Press, Tucson, pp. 75-95, 1993.
17. Bazan, G., Brown, L.E., Clayton, D.D., El Eid, M.F., Hartmann, D.H., and Truran, J.W., "Nucleosynthesis of  $^{26}\text{Al}$  in AGB Stars and Wolf-Rayet Stars," in Compton Gamma-Ray Observatory, AIP Conference Proceedings 280, pp. 47-51, 1993.
18. Truran, J.W. and Starrfield, S., "Gamma Rays and X Rays from Classical Novae," in Compton Gamma-Ray Observatory, Friedlander, M., Gehrels, N., and Macomb, D. (Eds.), AIP Conference Proceedings 280, pp. 163-167, 1993.
19. Starrfield, S., Shore, S.N., Sonneborn, G., Gonzalez-Riestra, R., Sparks, W.M., Truran, J.W., Dopita, M.A., and Williams, R.E., "The Production of  $^{22}\text{Na}$  and  $^{26}\text{Al}$  in Nova Explosions: The Observational Picture," in Compton Gamma-Ray Observatory, Friedlander, M., Gehrels, N., and Macomb, D. (Eds.), AIP Conference Proceedings 280, pp. 168-175, 1993.
20. Truran, J.W., Hayes, J., Livio, M., and Shankar, A., "On the Nature of the Super-Eddington Luminosities of Classical Novae," Preprint, 1993.
21. Starrfield, S., Truran, J.W., Sparks, W.M., Politano, M., Nofar, I., and Shaviv, G., "Nucleosynthesis in Nova Explosions," in Origin and Evolution of the Elements, Prantzos, N., Vangioni-Flam, E., and Casse, M. (Eds.), Cambridge University Press, Cambridge, pp. 337-343, 1993.
22. Starrfield, S., Truran, J.W. *et al.*, "Estimates of the  $^{22}\text{Na}$  and  $^{26}\text{Al}$  Production and Gamma-Ray Emission from Nova Cygni 1992 and Nova Herculis 1991," in "Cataclysmic Variables and Related Physics: 2nd Haifa-Technion Workshop on Cataclysmic Variable Stars," Regev, O. and Shaviv, G. (Eds.), Institute of Physics, Bristol, pp. 320-330, 1993.
23. Orio, M., Ögelman, H., Pietsh, W., Bianchini, M., Della Valle, M., Krautter, J., and Starrfield, S., "ROSAT All-Sky Survey Detections of Classical Novae and Related Objects," in Proceedings Cospar Conference, 1993.
24. Hollowell, D., Sparks, W.M., and Starrfield, S., "A Prediction of Compton Scattered Gamma-Rays from  $^{22}\text{Na}$  in Nova Cygni 1992," in "Cataclysmic Variables and Related Physics: 2nd Haifa-Technion Workshop on Cataclysmic Variable Stars," Regev, O. and Shaviv, G. (Eds.), Institute of Physics, Bristol, p. 290, 1993.
25. Shore, S.N., Sonneborn, G., Starrfield, S., Gonzalez-Riestra, R., and Ake, T.B., "The Early Ultraviolet Spectral Evolution of Nova Cygni 1992," Astronomical Journal, Vol. 106, p. 2408, 1993.
26. Shore, S.N., Sonneborn, G., Starrfield, S., Gonzalez-Riestra, R., and Polidan, R., "The Early Bolometric Evolution of Nova Cygni 1992," Astrophysical Journal, Vol. 421, pp. 344-349, 1994.
27. Hauschildt, P.H., Starrfield, S., Austin, S., Wagner, R.M., Shore, S.N., and Sonneborn, G., "Non-LTE Model Atmosphere Analysis of Nova Cygni 1992," Astrophysical Journal, Vol. 422, pp. 831-844, 1994.
28. Sion, E.M. and Starrfield, S., "Evolutionary Sequences of Very Hot, Low Mass, Accreting White Dwarfs," Astrophysical Journal, Vol. 421, pp. 261-268, 1994.

29. Bazan, G., Brown, L.E., Clayton, D.D., and Truran, J.W., " $^{26}\text{Al}$  Production in AGB Stars and 1.809 MeV Line Emission in the Galaxy," Preprint, 1994
30. Livio, M. and Truran, J.W., "On the Interpretation and Implications of Nova Abundances," Astrophysical Journal, in press, 1994.
31. Politano, M., Starrfield, S., Truran, J.W., Sparks, W.M., and Weiss, A., "Nucleosynthesis in Classical Nova Explosions on ONeMg White Dwarfs," Astrophysical Journal, in press, 1994.
32. Hayes, J. and Truran, J.W., "Powering of the Early Light Curves of the Classical Novae," Astrophysical Journal, in press, 1994.
33. Kashyap, V., Rosner, R., Schramm, D.N., and Truran, J.W., "Machos and the Diffuse X-Ray Background," submitted to Astrophysical Journal, 1994.
34. Truran, J.W., Glasner, S.A., and Hayes, J., "On the Nature of the Soft x-Ray Emission from Nova GQ Muscae 1983," Preprint, 1994.
35. Bazan, G. and Truran, J.W., "g-Process Nucleosynthesis in Asymptotic Giant Branch Stars as a Function of Initial Mass and Metallicity," Preprint, 1994.
36. Glasner, S.A. and Truran, J.W., "Soft X-Ray Outbursts on White Dwarfs," Preprint, 1994.
37. Fushiki, I., Thielemann, F.-K., and Truran, J.W., "General Implementation of Screening Effects in Thermonuclear Reaction Rates," Preprint, 1994.
38. Truran, J.W. and Livio, M., "Circumstantial Evidence for Significant Common-Envelope-Induced Mass Loss in Classical Novae Outbursts," in preparation, 1994.